



Hydrolyzed Collagen

Ready to support your patient's wound healing process

Some Patients are at a Higher Risk for Developing Surgical Wound Complications¹

Factors that increase these risks include:



Type I Collagen properties that support surgical wound progression²





CHEMOTACTIC




ADHESIVE

The Hydrolyzed Collagen Difference³

NATIVE COLLAGEN

HYDROLYZED COLLAGEN





Collagen wound into triple helix

Collagen fragmented into amino acids and peptides

2-5 days to degrade in a healthy patient

Ready for the body to use immediately

Insoluble

Soluble

Large molecule size

Small molecule size

High molecular weight

Low molecular weight



CellerateRX® Surgical Powder

A Type I bovine hydrolyzed collagen that aids in the management of surgical wounds

Indications

- Surgical Wounds
- Traumatic Wounds
- Partial and Full Thickness Wounds
- First and second- degree burns


The Benefits of CellerateRX Surgical Powder

- Provides the benefits of hydrolyzed collagen to the wound bed
- Hydrolyzed collagen fragments do not have to be broken down by the body before use
- Sterilized and packaged for use in the operating room
- Compatible with Negative Pressure Wound Therapy (NPWT)
- · Three-year shelf life

Application Process



Appropriate for Many Surgical Wounds







Cover the surgical wound as per your protocol

CellerateRX Surgical Ordering Information

PRODUCT#	PRODUCT DESCRIPTION	SIZE	
WCI-01-SACRXP	CellerateRX Surgical Powder	1g	
WCI-02-SACRXP WCI-05-SACRXP	CellerateRX Surgical Powder CellerateRX Surgical Powder	2g 5g	

Please contact us for more product information

800.205.7719 | sanaramedtech.com





© 2024 Sanara MedTech Inc. All rights reserved. CellerateRX and Activated Collagen are registered trademarks of Sanara MedTech Inc. MKT.01.32 Rev# 250819

CellerateRX Surgical Published Clinical Studies

JOURNAL	STUDY TITLE
Journal of Surgery	Activated Collagen Powder Significantly Reduces Surgical Site Infections in Patients Undergoing Elective Surgery ⁴
Orthopedics	The Effects of Platelet Rich Plasma and Activated Collagen on Wound Healing in Primary Total Joint Arthroplasty ⁵
JSM Neurosurgery and Spine	The Use of Sterile Bovine Type 1 Hydrolyzed Collagen to Support Surgical Wound Management: A Case Series ⁶
JSM Neurosurgery and Spine	Operative Closure Technique Utilizing Bovine Collagen Fragments in a Prospective Analysis of 102 Consecutive Neurosurgery Patients ⁷
JSM Neurosurgery and Spine	Retrospective Study to Evaluate the Use of Type 1 Bovine Hydrolyzed Collagen to Support Surgical Wound Healing After Spinal Surgery ⁸

References: 1. Cheadle WG. Risk factors for surgical site infection. Surg Infect (Larchmt). 2006;7 Suppl 1:S7-11. doi: 10.1089/sur.2006.7.s1-7. PMID: 16834549. 2. Mathew-Steiner SS, Roy S, Sen CK. Collagen in Wound Healing. Bioengineering (Basel). 2021 May 11;8(5):63. doi: 10.3390/bioengineering8050063. PMID: 34064689; PMCID: PMC8151502. 3. León-López A, Morales- Peñaloza A, Martínez-Juárez VM, Vargas-Torres A, Zeugolis DI, Aguirre-Álvarez G. Hydrolyzed Collagen-Sources and Applications. Molecules. 2019 Nov 7;24(22):4031. doi: 10.3390/molecules24224031. PMID: 31703345; PMCID: PMC6891674. 4. Nowrouzi R, Awad SS (2023) Activated Collagen Powder Significantly Reduces Surgical Site Infections in leman to Human Primary Total Joint Arthroplasty. Orthopedics. 2018; 41(2):e262-e267 6. Gitele Bovine Type 1 Hydrolyzed Collagen to Support Surgical Wound Management: A Case Series."JSM Neurosurg Spine08/22 7. Dickerman R, Ashley SRNP, Winters K. "Operative Closure Technique Utilizing Bovine Collagen Fragments in a Prospective Study to Evaluate the Use of Type 1 Bovine Hydrolyzed Collagen to Support Surgical Wound Healing After Spinal Surgery" JSM Neurosurg Spine 8(1):108.